

Assessing the effects of genotype-by-environment interaction on epigenetic, transcriptomic, and phenotypic response in a Pacific salmon

Kris A. Christensen ^{1,2,†}, Jérémy Le Luyer,^{3,†,‡} Michelle T. T. Chan,^{1,4} Eric B. Rondeau,^{1,2} Ben F. Koop,² Louis Bematchez ³, and Robert H. Devlin^{1,*}

¹Fisheries and Oceans Canada, West Vancouver, BC V7V 1N6, Canada

²Department of Biology, University of Victoria, Victoria, BC V8P 5C2, Canada

³Département de Biologie, Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, QC G1V 0A6, Canada

⁴Molecular Biology and Biochemistry Department, Simon Fraser University, Burnaby, BC V5A 1S6, Canada

[†]Equal contribution.

[‡]Present address: IFREMER, EIO UMR 241, Labex CORAIL, Unité RMPF, Centre Océanologique du Pacifique, Vairao – BP 49 Vairao, Tahiti, Polynésie française.

*Corresponding author: Fisheries and Oceans Canada, 4160 Marine Drive, West Vancouver, BC V7V 1N6, Canada. robert.devlin@dfo-mpo.gc.ca

Abstract

Genotype-by-environment (GxE) interactions are non-parallel reaction norms among individuals with different genotypes in response to different environmental conditions. GxE interactions are an extension of phenotypic plasticity and consequently studying such interactions improves our ability to predict effects of different environments on phenotype as well as the fitness of genetically distinct organisms and their capacity to interact with ecosystems. Growth hormone transgenic coho salmon grow much faster than non-transgenics when raised in tank environments, but show little difference in growth when reared in nature-like streams. We used this model system to evaluate potential mechanisms underlying this growth rate GxE interaction, performing RNA-seq to measure gene transcription and whole-genome bisulfite sequencing to measure gene methylation in liver tissue. Gene ontology (GO) term analysis revealed stress as an important biological process potentially influencing growth rate GxE interactions. While few genes with transcription differences also had methylation differences, in promoter or gene regions, many genes were differentially methylated between tank and stream environments. A GO term analysis of differentially methylated genes between tank and stream environments revealed increased methylation in the stream environment of more than 95% of the differentially methylated genes, many with biological processes unrelated to liver function. The lower nutritional condition of the stream environment may cause increased negative regulation of genes less vital for liver tissue function than when fish are reared in tanks with unlimited food availability. These data show a large effect of rearing environment both on gene expression and methylation, but it is less clear that the detected epigenetic marks are responsible for the observed altered growth and physiological responses.

Keywords: epigenetics; transcriptome; GxE; transgenic; growth hormone; salmonid

Introduction

Phenotypic plasticity is a characteristic of species or strains in which phenotype (e.g., morphology, physiology, and behavior) is not fixed between environments. Because phenotypic plasticity may confer a fitness advantage by allowing non-genetic modification of traits, its role in modifying evolutionary trajectories through genetic accommodation or assimilation has been a fascinating question in evolution and remains a matter of ongoing discussion (Yeh and Price 2004; Laland et al. 2014; Schlichting and Wund 2014; Levis and Pfennig 2016; Charlesworth et al. 2017; Oostra et al. 2018). Phenotypic plasticity may provide higher adaptive potential than the accumulation of *de novo* mutations due to its ability to affect multiple individuals rapidly, respond to environment-specific triggers, and harness existing (yet cryptic) genetic variation (Levis and Pfennig 2016). For instance, field studies in the threespine stickleback (*Gasterosteus aculeatus*)

demonstrated the potential of phenotypic plasticity in promoting adaptive radiation of two fish ecotypes through parallel selection of developmental phenotypes (Wund et al. 2008).

Genotype-by-environment (GxE) interaction is a type of phenotypic plasticity influenced by both genotype and environment and where the phenotypic responses of organisms with one genotype are different from those of another genotype between two or more environments—specifically, in a non-parallel fashion. It is thought that gains in selective breeding programs can be hampered when unexpected GxE interactions are not included in genetic evaluations (Mulder 2016). Similarly, in risk assessment of transgenic salmon, GxE interactions [e.g., physiology (Löhmus et al. 2010), predation (Sundström et al. 2007), growth, survival (Vandersteen et al. 2019), and antipredator behavior (Sundström et al. 2016)] might lead to mistaken conclusions if non-transgenic and transgenic salmon have non-parallel reaction norms across

Received: October 28, 2020. Accepted: January 13, 2021

© Crown copyright 2021.

This article contains public sector information licensed under the Open Government Licence v3.0 (<http://www.nationalarchives.gov.uk/doc/open-government-licence/version/3/>)

environments (Devlin *et al.* 2015). While commercial use of transgenic fish has historically been of concern, transgenic Atlantic salmon (*Salmo salar*) are now being produced commercially (Waltz 2017) in highly secure land-based culture facilities. If such measures are not applied to the rearing of transgenic fish, these fish may gain access to natural environments. Identifying ecological effects of such an outcome is complicated by GxE interactions affecting phenotype.

The underlying mechanisms of phenotypic plasticity and GxE interactions are of interest as they may provide insight into how the environment interacts with the genome and allows improved predictions of the effects of different environments on phenotype (Nicotra *et al.* 2010). Many studies have focused on the epigenetic mechanisms underlying phenotypic plasticity [e.g., drought tolerance (Herman and Sultan 2016), fungus in multiple environments (Kronholm *et al.* 2016), migration (Baerwald *et al.* 2016), growth in various sugar concentrations (Herrera *et al.* 2012), and effect of rearing environment (Luyer *et al.* 2017)]. Potentially other mechanisms influencing gene expression could underlie phenotypic plasticity and GxE interactions (Smith and Kruglyak 2008; Gutierrez *et al.* 2009).

Epigenetics is a broad field that can refer to multiple and varied pathways: from DNA methylation to self-regulating transcription factors (reviewed in Martin and Fry 2018; Cavalli and Heard 2019). Epigenetic gene regulation differs from general gene regulation by duration, where effects might extend long after the initial signal has faded (see Cavalli and Heard 2019 for other definitions). Plasticity is related to epigenetics in that it is the ability of regulatory mechanisms (e.g., transcription factors) to be influenced by environmental differences to change or reverse epigenetic factors already in place and governing cellular characteristics (Cavalli and Heard 2019).

For environmentally induced epigenetic changes, DNA methylation is the best known and studied epigenetic marker (Martin and Fry 2018). Methylation occurs when a methyl group is added to a cytosine by a DNA methyltransferase that has been triggered by environmental signals (e.g., chemicals interacting with DNA methyltransferases, availability of the methyl donor SAM, or the availability of cytosines due to transcription factor binding or the presence of trimethylated histone H3 Lys4 near specific genes; Martin and Fry 2018; Greenberg and Bourc'his 2019). In humans, methylation appears to play a role in regulating genes like insulin-like growth factor 2 (i.e., less DNA methylation after exposure to famine) and other genes depending on environmental exposure (Heijmans *et al.* 2008; Martin and Fry 2018). DNA methylation has typically been associated with transcriptional repression by decreasing transcription factor binding, increasing heterochromatin formation, and by recruiting proteins that silence gene transcription (Greenberg and Bourc'his 2019). DNA methylation is likely more complex than this simple model. For example, in human phagocytic cells, gene transcription appeared to be independent of DNA methylation (Pacis *et al.* 2019). Although this may be due to detection limits and the statistics used in identifying differentially methylated regions (including which methylation mark is being monitored) (Pacis *et al.* 2019), it is also likely gene expression changes are regulated through many pathways.

Transgenic salmon have dramatically faster growth due to the integration of the growth hormone transgene construct at a single genomic locus (Devlin *et al.* 1994; Phillips and Devlin 2010). In the wild, non-transgenic salmonids possess two growth hormone genes (GH1 and GH2) that influence many biological processes (e.g., growth, osmoregulation, and immune function) and are

central to a complex signaling system that integrates information from multiple sources (Björnsson *et al.* 2002). Central figures in relaying GH signaling in salmonids are the GH-binding proteins (GHBP; regulates circulating GH), growth hormone receptor (GHR; highest density in the liver), and insulin-like growth factor 1 (IGF1; gene expression stimulated by GH and primarily produced in the liver) (Björnsson *et al.* 2002). The transgenic coho salmon in this study have an extra and constitutively active versions of the growth hormone gene (Devlin *et al.* 1994) causing growth hormone transgene to be expressed in all tissues and is known to have strong effects on gene expression, growth, and physiology (Rise *et al.* 2006; Devlin *et al.* 2009, 2015).

In previous studies (Sundström *et al.* 2007, 2016; Vandersteen *et al.* 2019), GxE interactions were observed for various traits when non-transgenic and transgenic coho salmon were reared both in stream and in tank environments. Specifically, the elevated growth of transgenic individuals seen in simple tank environments was not observed in the naturalized and complex stream environment where non-transgenic and transgenic coho salmon grew at the same rate (Sundström *et al.* 2007). The plasticity and GxE interaction effects observed in GH transgenic salmon provide a useful model system to better understand epigenetic mechanisms affecting phenotype in vertebrates. We sought to identify underlying gene expression differences related to GxE interactions and to assess if gene expression GxE interactions were influenced by differential methylation.

Materials and methods

Full details of the materials and methods used in this study can be found in File S1. Briefly, growth hormone transgenic and non-transgenic coho salmon (Sundström *et al.* 2007) were reared in tank or simulated stream environments. Liver tissues were collected from two sets of six individuals from each environment and genotype and each group of six was pooled as a single replicate (two replicates for each environment and genotype). RNA-seq and whole-genome bisulfite sequencing was performed on each replicate (NCBI SRA accessions: SRX9236012–SRX9236027). The analyses of the RNA-seq and whole-genome bisulfite sequencing data included pairwise comparisons and a genotype-by-environment analysis. Methylation data was analyzed based on promoter region, gene body region, and intergenic sites (scripts can be found in File S2). Gene ontology analyses were used to identify enriched categories of genes.

Data availability

Sequence data were deposited to the NCBI's sequence read archive available through the BioProject ID: PRJNA667073, and R/python scripts are available as supplementary files (File S2—GSA figshare). All supplemental files can be accessed through figshare: <https://doi.org/10.25387/g3.13542158>.

Results

Body weight and specific growth rate

Body weights did not reflect growth rate differences among environments or genotypes as salmon groups were size-matched to avoid confounding effects of developmental stage which occurs when analyzing strains with different growth rates sampled at the same age (Figure 1A). A significant interaction between environment and genotype ($P < 2e-16$) for specific growth rate (SGR) was identified using a two-way ANOVA test (Figure 1A). The average non-transgenic SGR was increased relative to the average




Figure 1 Genotype-by-environment influence on weight and specific growth rate of coho salmon. (A) Average coho salmon size per group were matched to avoid confounding effects of developmental stage. No variables were significantly different for weight. (B) The specific growth rate for each group is shown (mean \pm SEM and individual specific growth rate). Environment, genotype, and the interaction of both were all significant with $P < 2e-16$, $P = 8.59e-15$, and $P < 2e-16$, respectively.




Figure 2 Multidimensional scaling plot (MDS) of gene expression data and a Venn diagram of overlapping differentially expressed genes (DEGs). (A) A MDS plot of transcript counts for all genes expressed in the liver of transgenic and non-transgenic coho salmon reared in either stream or tank environments (as expressed in log fold change, logFC). Each point in the MDS plots represent six pooled individuals. (B) Overlapping DEGs between comparisons.

transgenic SGR in the stream environment but decreased in the tank environment (Figure 1B).

RNA-seq and WGBS sequence data

RNA-seq paired-end counts ranged from 36.6M to 59.2M per pooled sample (NCBI BioProject: PRJNA667073). After trimming, a mean of 42.9 ± 2.5 M reads were retained per pooled sample. Read alignment rates averaged $89 \pm 3.8\%$ for properly-paired reads mapping to the coho reference genome. WGBS paired-end sequence counts ranged from 481.9M to 512.5M per pooled sample. After trimming, a mean of 441.6 ± 4.1 M reads were retained per pooled sample. Overall, mapping rates were $45.8 \pm 0.23\%$ for uniquely and paired reads [around $17\times$ coverage assuming an estimated 2.4 Gbp genome, which is likely a higher coverage than necessary for analysis (Ziller et al. 2015)].

RNA-seq analysis

Transcription of genes in the liver was greatly influenced by the rearing environment (Figure 2A, Table 1). Replicate groups clustered closely indicating shared responses to both genotype and environment. Only 382 DEGs were identified between transgenic

and non-transgenic fish in the stream environment compared to 3134 differentially expressed genes (DEGs) in a tank environment (Figure 2B, Table 1). This difference between environments was consistent with the growth rate analysis, where transgenic and non-transgenic fish had similar SGRs in a stream environment, but more divergent SGRs in a tank environment (Figure 1A). Three enriched biological process GO terms were identified between transgenic and non-transgenic salmon in the stream environment: *complement activation*, *complement activation, alternative pathway*, and *organic hydroxy compound catabolic process*. In the tank environment, 304 biological process enriched GO terms were identified (Supplementary File S3, Figure S1), many of which were shared with GxE enriched gene categories (see below). There were 759 shared DEGs between this comparison and the GxE comparison (84% of the GxE DEGs—see below).

Both non-transgenic and transgenic salmon had a similar number of liver DEGs between the stream and tank environments but only about 50% of those DEG were shared between both groups (Figure 2B, Table 1, non-transgenic: 5826, transgenic: 6216, shared: 3083). There were 779 enriched biological process GO terms identified between environments for non-transgenic

salmon and 767 for transgenic salmon, with 452 shared (Supplementary File S3, Figure S2). Major themes from both sets of GO terms include: metabolism, ribosome biogenesis, stress, immune system, cell death, chromatin organization, and gene expression regulation (including epigenetic regulation).

From the ANOVA-style analysis (any gene that was differentially expressed between any of the comparisons), 9615 DEGs were identified and 901 DEGs were classified from the GxE analysis (Figure 3A, Supplementary File S3). There were 311 enriched biological processes, 7 cellular components, and 37 molecular function GO terms identified from the GxE analysis (Figure 3B, Supplementary File S3). The GO term with the lowest P-value was *response to hormone* and is consistent with the growth hormone transgene physiological alteration observed in SGR (Sundström et al. 2007). Many of the other lowest P-value enriched GO terms were responses to various stimuli (e.g., lipid, insulin, stress, and nutrient levels), signaling (e.g., insulin receptor signaling pathway, JAK pathway signal transduction adaptor activity, and interleukin-7-mediated signaling pathway), and regulation (e.g., regulation of primary metabolic process, regulation of complement activation, and regulation of hormone secretion) were other main themes found in these GO terms (File S3).

Table 1 Number of differentially expressed genes

	Stream Non- transgenic	Tank Non- transgenic	Tank Transgenic
Stream Non-transgenic	NA	5826	<i>a</i>
Stream Transgenic	382	<i>a</i>	6216
Tank Transgenic	<i>a</i>	3134	NA

^a Not-tested.




Figure 3 Genotype-by-environment interactions of liver mRNA transcription in a stream and tank environment. (A) A heatmap of transcript counts per million (CPM) values of 901 DEGs with GxE interactions. The NT symbol represents non-transgenic and T represents transgenic. For each experimental condition there were two replicates of pooled individuals. (B) Enriched GO terms of the genes with GxE interactions after complexity reduction [i.e., cluster representatives are shown (Supek et al. 2011)] of the 311 biological process GO terms originally identified. The x and y-axes are coordinates from multidimensional scaling of GO term semantic similarities into two dimensions. The size and color of each GO term bubble reflects the P-value, with larger and blue bubbles with the lowest P-values.

To better understand GxE transcription interactions, the 901 GxE DEGs were clustered based on transcript expression patterns (Figure 4, A and B). Out of 24 identified clusters, enriched GO terms could be classified for 11 of them (Figure 4C). For clusters 1-3, transgenic transcript expression tended to increase from a stream to tank environment, while expression decreased for non-transgenic fish between environments (Figure 4C). The lowest P-value categories for each of these clusters were *positive regulation of vitamin D receptor signaling pathway*, *regulation of cellular metabolic process*, and *negative regulation of interleukin-2 secretion* in increasing order (File S3).

The GO category with the lowest P-value from cluster 8, *ribosome biogenesis*, is suggestive that transcription of genes involved in ribosome biogenesis increases in both transgenic and non-transgenic salmon in tank environments, but to a greater extent in transgenic fish (Figure 4C, Supplementary File S3). This pattern of transcription mirrors SGR between environments. Transgenic coho salmon have decreased transcription of genes related to clusters 9–10 in a tank environment. GO categories with the lowest P-value from these clusters include, in increasing cluster order, *digestion* and *complement activation* (File S3). Transgenic fish displayed an opposite transcription pattern for these categories. Transcription in cluster 11 increases from stream to tank environments in transgenic salmon and decreases in non-transgenic salmon (similar to clusters 1–3). The top enriched GO category for this cluster was *regulation of cardiac chamber formation*.

Clusters 13, 21, and 22 all have similar gene transcription patterns with increased transcription in a tank environment for non-transgenic fish and decreased for transgenic fish. Top enriched GO terms for these clusters were *acute inflammatory response to antigenic stimulus*, *positive regulation of cell death*, and *glucose metabolic process*, respectively (File S3). Hepatic gene transcription decreased for genes in cluster 20 in a tank environment for transgenic and non-transgenic salmon, but the drop in expression was




Figure 4 Genotype-by-environment interactions of liver mRNA-transcription clustering. (A) A heatmap of Spearman correlation coefficients with genes clustered with similar gene transcription. Each of these genes had transcript GxE interaction. The NT symbol represents non-transgenic and the T represents transgenic. (B) Two example clusters are shown of the 24 identified. Colors of the heatmap are based on transcript counts per million (CPM) values. (C) Illustrations of expression patterns for each cluster found to have enriched GO terms (does not reflect actual expression values). Some clusters were grouped in this figure, but each were individually analyzed for GO term enrichment. Boxes above graphs show summaries of enriched GO terms found in clusters. Blue boxes show GO term summaries related to metabolism, orange boxes related to the immune system, and green boxes related to vitamin D. Other box colors were unique categories.

greater for non-transgenic salmon. The top enriched GO term for this cluster was *response to cesium ions* (File S3).

WGBS analysis

Methylation patterns differed between both promoter regions and intergenic sites compared to gene regions (Figure 5, Supplementary Figures S3 and S4). As was the case for RNA-seq analysis, replicate groups clustered closely. Most of the changes in hepatic promoter methylation and intergenic site methylation occurred between transgenic and non-transgenic salmon in both stream and tank environments (Figure 5B). There were 4 out of 382 DEGs (1%) that were also differentially methylated between non-transgenic and transgenic salmon in the stream environment and 34 out of 3134 DEGs (1%) in the tank environment. Methylation at 18 promoter regions displayed GxE interaction (Table 2, Supplementary Figure S5). In total, 823 promoter regions were differentially methylated in the ANOVA-style analysis (differentially methylated in any of the comparisons). No enriched GO terms were identified for any promoter methylation comparison. GxE methylation patterns were identified at 485 intergenic sites, and 76050 intergenic sites were found to be differentially methylated in the ANOVA-style analysis (File S4).

In gene regions, most changes in methylation were observed between environments rather than between genotypes as seen in promoter regions and intergenic sites (Figure 5B). More than 95% of the observed changes were from increased methylation of genes in the stream environment. Out of the 6216 DEGs found between environments for transgenic salmon, 221 (3.6%) were also

differentially methylated, while 222 out of the 5826 DEGs (3.8%) for the non-transgenic salmon were differentially methylated. Common enriched GO terms found between stream and tank environments for both transgenic and non-transgenic comparisons include: *behavior*, *locomotion*, *nervous system development*, and *homophillic cell adhesion via plasma membrane adhesion molecules* (Figure 6, Supplementary File S4). Nine GO terms from these two comparisons were shared with those found in the GxE gene expression analysis, but these terms were from broad categories (e.g., *cell differentiation*, *cellular developmental process*, and *developmental process*). There were 63 gene regions with GxE interactions (File S4). No enriched GxE GO terms were identified, but there were several important transcription factors influencing growth identified by this analysis including: neurogenic differentiation factor 1-like, transcription factor jun-B-like, and cyclic AMP-dependent transcription factor ATF-1-like (Supplementary Figure S6). Only the neurogenic differentiation factor 1-like gene had methylation patterns expected to influence gene expression (i.e., an increase in methylation corresponded to a decrease in gene expression between environments and a decrease in methylation corresponded to an increase in gene expression) (Supplementary Figure S6).

Discussion

The goal of this study was to better understand how vertebrate phenotypes can be affected by genotype and environment and their interactions at a genomic level. We have utilized a GH




Figure 5 Differentially methylation (DM) site summary information. (A) Multidimensional scaling (MDS) plots of methylation data ($\log_2(\text{Methylated_counts} + 2) - \log_2(\text{Unmethylated_counts} + 2)$) for all regions or sites in the liver of transgenic and non-transgenic coho salmon reared in either stream or tank environments (expressed in log fold change, logFC). The first MDS plot was generated from all promoter regions, the second was generated from all gene regions, and the last from all methylation sites outside of gene and promoter regions. (B) Venn diagrams of overlapping promoters, genes, or intergenic sites found between different comparisons. Below the promoter and gene Venn diagrams are lines showing the number of overlapping DM regions between promoter and gene based analyses.

Table 2 Genes with GxE methylation interactions in the promoter region

NCBI Accession	Description
LOC109904709	secretin receptor-like
sass6	spindle assembly abnormal protein 6 homolog
LOC109889987	prolyl 4-hydroxylase subunit alpha-1-like
ube2j2	LOW QUALITY PROTEIN: ubiquitin-conjugating enzyme E2 J2
LOC109899332	transmembrane protein 14C-like
rbfox3	RNA binding protein fox-1 homolog 3
LOC109901502	thyrotropin-releasing hormone receptor-like
LOC109901621	uncharacterized protein
LOC109904333	zinc finger protein 271-like
LOC109905467	DNA-directed RNA polymerase I subunit RPA34-like
LOC109905896	amyloid protein-binding protein 2-like
LOC109909977	patched domain-containing protein 3-like
setd6	N-lysine methyltransferase SETD6
LOC109876066	sodium/calcium exchanger 3-like
LOC109879198	vesicle transport protein SEC20-like
LOC109882311	flocculation protein FLO11-like
LOC109884608	glutamate receptor ionotropic%2C NMDA 2A-like
LOC109885425	kelch domain-containing protein 1-like

transgenic salmon model where the genetically modified salmon grow much quicker than non-transgenic salmon in a tank environment but grow similarly in a stream environment (Sundström et al. 2007). From the RNA-seq data, we identified genes with genotype, environmental, and GxE interaction transcription

patterns in the liver. Clustering genes by GxE interaction helped us to identify which biological processes changed in a similar fashion as seen for growth. To better understand how genes with GxE interactions were regulated, whole-genome bisulfite sequencing was performed on the same samples and tissue. Although we observed that many genes had increased methylation in the stream environment compared to the tank environment and reduced expression for those with increased methylation overall, we found little overlap with changes in methylation and changes in gene expression. Each of these observations are described in more detail below.

Specific growth rate

Both non-transgenic and transgenic salmon grew faster in the tank environment than in a stream environment. Environmental, physiological, and behavioral factors likely contribute to the two-fold increase in SGR in tanks [e.g., availability of food, nutritional capabilities, and suppression of predator avoidance (Devlin et al. 2015)]. To uncover which genes influence or underlie these responses, we searched for genes with liver transcription profiles with complex responses including GxE interactions.

RNA-seq genotype differences

In the stream environment, only three enriched GO terms (complement activation; complement activation, alternative pathway; and organic hydroxy compound catabolic process among 382 DEGs) were identified from the comparison between non-transgenic and transgenic salmon liver tissue. Two of these were related to complement activation, suggesting that some




Figure 6 Enriched GO terms identified from genes with differential methylation between stream and tank environments. GO terms complexity was reduced using Revigo software (Supek et al. 2011), and GO terms with labels were chosen to maximize the shared GO terms and based on ease of interpretation or *P*-value. The x and y-axes are coordinates from multidimensional scaling of GO term semantic similarities into two dimensions. (A) Enriched GO terms identified between the stream and tank non-transgenic comparison of gene body methylation. (B) Enriched GO terms identified between the stream and tank transgenic comparison of gene body methylation.

differences between liver transcription of transgenic and non-transgenic salmon in the stream environment could be due to environment specific pathogens (Thorgersen et al. 2019). In the tank environment, differences in B cell and T cell regulation were noted between transgenic and non-transgenic salmon, but not with the complement activation GO term. As these immune-related differences were specific to only one of the environments, they represent GxE interactions and were also detected in the GxE analysis.

Recent studies found that GH transgenic salmon have altered immune responses to pathogen mimics in tank environments (Alzaid et al. 2018; Kim et al. 2019). Taken together with the current study, transgenic, and non-transgenic salmon appear to respond differently to pathogen exposure in multiple environments. In the current study, the transgene appeared to influence immune system differences while the environment appeared to influence which aspect of the immune-system (complement activation vs T cell or B cell regulation) was different between transgenic and non-transgenic salmon. The environmental differences could be driven by different pathogen communities, but it could also be as simple as temperature differences (Ignatz et al. 2020). Alzaid et al. (2018) hypothesized that the immune system may modulate growth as a trade-off. Further research into these immune differences may improve cultivation practices of farmed salmon and risk assessments of transgenic salmon as immune function is important to both objectives.

Many of the enriched GO terms identified in the comparison between transgenic and non-transgenic coho salmon in the tank environment were also shared with the GxE comparison (115 shared out of the 304 in the tank environment). Most of the remaining GO terms, without exact matches, shared similar biological functions (e.g., regulation of mitotic cell cycle vs regulation of mitotic cell cycle phase transition). For this reason, discussion of these GO terms is reserved for the GxE section below. It is important to note here, however, that these differences were not seen in the stream environment. This suggests that controlled studies in tank environments may not accurately represent results relevant to stream or wild environments, and, as such,

the best risk assessments would incorporate data from multiple environments.

RNA-seq environmental differences

An unanticipated finding from comparing stream and tank environments was that there might be an influence of food availability in the stream environment, as the *cellular response to starvation* enriched GO term was identified between environments for both the transgenic and non-transgenic salmon. This result was surprising because transgenic and non-transgenic salmon both had positive growth in stream and tank environments. Also, transgenic and non-transgenic salmon were fed to excess in both environments (adding more natural food to this experimental stream system does not increase growth rates; Vandersteent et al. 2019). We observed that in a stream environment both genotypes had reduced growth relative to the tank environment, but both environments still had positive growth overall. Some questions arise from these observations, e.g., are the fish in the stream environment responding specifically to reduced food supply, and are the cellular mechanisms initiated by lower levels of available food also activated by other conditions or under submaximal feed regimes where fish must compete for available natural prey?

There are environmental and behavioral differences that result in reduced nutrition between stream and tank environments [e.g., the nutrition of the prey in the stream environment may be different than feed pellets causing decreased growth rate (Holm et al. 1984; Vandersteent et al. 2019)]. It is possible that submaximal feed levels in the stream environment initiated cellular responses similar to starvation if these are more general purpose cellular responses and do not strictly respond to complete nutrient deprivation. This explanation fits with the observation of reduced growth in the stream environment, the *cellular responses to starvation* GO term, and the feeding regime used in this study.

Many other shared differences of non-transgenic and transgenic salmon were observed between stream and tank environments. They include enriched GO categories related to circadian rhythm, metabolism, protein folding, autophagy, cellular response to stress, chromatin silencing, demethylation, and

regulation of gene expression (epigenetic) among many others. While these GO terms may be informative regarding trait differences between fish reared in tank and enriched stream environments, they do not offer obvious insight into growth (the trait under investigation).

RNA-seq GxE interactions

From the 901 DEGs with GxE transcription interactions in the liver and corresponding enriched GO terms, several inferences can be made regarding the physiological and gene transcription responses regulating the SGR GxE interaction. First, the enriched GO term with the lowest *P*-value, *response to hormone*, supports that the growth hormone transgene was a main genetic influence in the SGR GxE interaction. Other enriched GO terms further support the influence of the growth hormone in the SGR GxE interaction including: *JAK pathway signal transduction adaptor activity*, *ERK1 and ERK2 cascade*, and *insulin receptor signaling pathway* (insulin-like growth factor I was one of the 901 GxE DEGs) as expected from the known effects of growth hormone (Dehkhoda et al. 2018). This result was expected since salmon with similar genetic backgrounds were used—with the exception of the transgene. Second, several other enriched GxE interaction enriched GO term categories (e.g., *metabolism*, *glycogen metabolism*, *gluconeogenesis*, *response to insulin*, *response to cAMP*, *response to glucocorticoid*, *negative regulation of growth*, and *response to nutrient levels*) implicate genes involved in nutrition as downstream drivers of the SGR GxE interaction. Several of these enriched GO terms were found within clusters 21 and 22, which helps us to better understand gene expression patterns associated with these GO terms. In clusters 21 and 22, gene transcription has an inverse relationship to SGR patterns between environments (i.e., gene transcription is lower in the stream environment and higher in the tank environment for the non-transgenic salmon compared to transgenic salmon). One explanation for these expression patterns and the enriched GO terms is that glycogen metabolism and gluconeogenesis may be occurring to a greater extent in liver tissues of transgenic salmon in a stream environment and in non-transgenic salmon in the tank environment.

In humans, glycogenolysis in the liver provides glucose to the rest of the body during short fasting periods and gluconeogenesis provides glucose to the rest of the body for longer starvation periods or with increased stress (Zhang et al. 2019). In the current study, growth was positive in both stream and tank environments, and with feed in excess in the tank environment, starvation seems unlikely (see above for discussion on starvation). The enriched GO terms *cellular response to glucocorticoid stimulus* and *cellular response to stress* in clusters 21 and 22 are indicative of a stress response, which may also promote gluconeogenesis and glycogen storage (Kuo et al. 2015). Stress may influence growth by reducing attempts to feed or by reduced feed absorption (Van Weerd and Komen 1998). Stress has also been shown to cause cortisol and glucose GxE interactions in other fish species (Van Weerd and Komen 1998). Stress responses may help explain the difference in growth data seen between transgenic and non-transgenic salmon from other studies under various conditions and in different environments (Vandersteen et al. 2019). Similarly, cortisol and glucose responses can vary based on stressor, genotype, and early life history (Van Weerd and Komen 1998) making stress a good candidate for the growth discrepancy observed in previous studies. We note that GH transgenic coho salmon do show altered carbohydrate (complex and glucose) metabolism, but GxE interactions have not been assessed in these studies (Higgs et al. 2009; Panserat et al. 2014).

With regards to the GxE enriched GO terms, liver protein synthesis has an association with SGR. The gene transcription patterns in cluster 8 mirror the SGR patterns, and this cluster of genes is enriched for genes involved in ribosome biogenesis. Growth and ribosome biogenesis is intimately linked as protein production is necessary for cell growth (Lempiäinen and Shore 2009).

From the enriched GO term and clustering analyses, we were able to link SGR GxE patterns to gene expression patterns of the genes downstream of the growth hormone, stress-related genes, and genes responsible for ribosomal biogenesis. We also identified several other categories of genes with GxE transcription interactions that may not have a direct influence on SGR, but may be important indirectly or for other unmeasured traits. These include genes with immune function (e.g., *complement activation*, *regulation of immune system process*, *response to interleukin-7*, and *defense response*), tissue regeneration (e.g. *animal organ regeneration* and *liver regeneration*), vitamin metabolism (e.g., *vitamin D3 metabolic process*, *vitamin D catabolic process*, and *response to vitamin B2*), and pigment formation (e.g., *pigment metabolic process*). Future research will be needed to better understand why these gene categories also showed GxE interactions.

WGBS genotype differences

While there were extensive differences between non-transgenic and transgenic methylation patterns in promoter regions and intergenic sites, there were no enriched GO terms identified for the promoter regions. There is likely important biological significance and possibly overlap with the SGR phenotype due to these differences, but the evidence suggests they underlie only a fraction of gene expression differences in the liver between transgenic and non-transgenic salmon. For example, gene transcription is markedly different between non-transgenic and transgenic salmon in the tank environment (3134 DEGs), but similar in a stream environment (382 DEGs). Meanwhile, there are similar levels of differentially methylated promoters and intergenic sites between non-transgenic and transgenic salmon in tank (366 differentially methylated promoters and 31,016 intergenic sites, File S5) and stream environments (326 differentially methylated promoters and 29,194 intergenic sites). While the overall increased methylation and reduced gene expression (see below) in the stream environment for these genes with increased methylation could also be responsible for the few DEGs in the stream environment, we did not observe a high proportion of overlapping DEGs between environments and genes that were differentially methylated (e.g., 221 differentially methylated gene regions shared out of 6216 DEGs between environments for transgenic salmon, File S4). One explanation for the divergence between liver gene transcription and methylation data is that the data reflects unobserved factors at the level of the organ or whole organism not accounted for with single tissue data. This hypothesis is supported by gene region methylation patterns seen between environments (see below).

WGBS environmental differences

Distinctive liver methylation patterns for gene regions were observed between stream and tank-reared salmon for both non-transgenic and transgenic salmon. Many of the enriched biological process GO terms between environments were shared between non-transgenic and transgenic salmon (341 non-transgenic, 215 transgenic, 154 shared). Surprisingly, the biological process GO terms between environments were mostly unrelated to known liver biological processes (e.g., *learning* or

memory, muscle contraction, and thigmotaxis). More than 95% of the gene regions involved in these biological processes had increased methylation in the stream environment. Because these GO terms are likely important for other organs rather than for liver tissues and the genes from these terms have increased methylation, we suggest that they are under strict gene transcription repression in the liver due to environmental factors [e.g., nutritional differences between prey and hatchery feed (Holm et al. 1984; Vandersteen et al. 2019)]. The average counts per million (CPM) value for all genes and samples after filtering was 38.4 ($n=26,084$, stdev: 493.4), whereas for the subset of genes found to have significantly increased methylation in a stream environment, the average CPM was 33.8 ($n=1262$, stdev: 523.1) for the non-transgenic comparison and 11.4 ($n=1250$, stdev: 104.3) for the transgenic comparison. In a previous study, hatchery coho salmon (analogous to tank-reared in this study) had increased methylation of most differentially methylated regions in muscle tissue compared to natural salmon (analogous to stream-reared) (Luyer et al. 2017). While the pattern of methylation was the opposite for this study, it is interesting that the trend for both tissues was unidirectional.

WGBS GxE interactions

There were 18 promoter regions and 63 gene regions identified with GxE methylation patterns. None of these promoter or gene regions overlapped with genes with GxE gene expression patterns, and only three promoter and ten gene regions with methylation GxE patterns were found to be differentially expressed in an ANOVA-style analysis of gene expression data. Of the 901 genes with GxE patterns, 23 promoters and 113 gene regions were found to have differential methylation in an ANOVA-style analysis. Both observations lead us to conclude that the majority of GxE transcription was not influenced directly by methylation and that other mechanisms influencing gene transcription were likely responsible. The influence of long-distance methylation and methylation of hub genes involved in large networks may still play important regulatory roles in the observed GxE transcription, but were not examined in this study. Another possibility not explored in this study, but may add additional insight is the influence of methylation position (e.g., could changing the position of methyl groups within the promoter region change gene expression while not appearing to be significantly different). As mentioned in the Introduction, gene expression and DNA methylation may not have a direct relationship, as exemplified in human phagocytes (Pacis et al. 2019), and subtle or complex relationships may not be accounted for in the type of analysis used. However, a disconnect between methylation and gene expression has also been observed in other studies, where gene expression differences were not associated with methylation differences for the majority of genes (Cunningham et al. 2019; Natri et al. 2020). These observations do not suggest that DNA methylation did not play an important role in gene expression regulation in general, but that the gene expression differences were not obviously linked to changes in methylation. For example, in the promoter region there was two clusters of genes (Supplementary Figure S3), one with a peak with around 25% average methylation and another with 90% average methylation. The first cluster had a peak at around 10 counts per million (CPM, measure of transcription), while the second peak was below 1 CPM. These clusters are consistent with expectations of decreased gene expression with increased promoter methylation and was seen in the majority of genes.

Methylation of DNA coding transcription factors may still play a significant role in the observed SGR GxE interaction. For example, two transcription factors (transcription factor jun-B-like and neurogenic differentiation factor 1-like) with GxE methylation patterns in gene regions may both play roles in growth (Andrali et al. 2008; Raffaello et al. 2010). Transcription factor jun-B plays an essential role in skeletal muscle mass maintenance, with increased expression associated with increased muscle mass and decreased expression with atrophy (Raffaello et al. 2010). Transcription of this factor is congruent with the SGR GxE interaction (Supplementary Figure S6), but did not reach significance in the GxE analysis. Methylation differences do not correlate with the expression pattern with this gene. Neurogenic differentiation factor 1-like regulates insulin gene expression in the pancreas and liver (Andrali et al. 2008). While this gene did not have significant GxE gene expression, it did have a gene expression pattern that was the opposite of the SGR GxE profile. If DNA methylation of gene or promoter regions in the liver underlies the SGR GxE profile, it likely does so through a gene that can regulate many other genes (i.e., a gene hub) like these transcription factors because there is little evidence for methylation differences that could explain the 901 GxE DEGs otherwise. More likely, DNA methylation and other regulators of gene expression play an elaborate role in the observed liver GxE gene expression profile.

Conclusions

Liver gene transcription data in transgenic and non-transgenic salmon pointed to genes related to stress as candidates underlying the GxE pattern of growth in this study. For the model system used here, if transgenic and non-transgenic coho salmon respond to environments differently in terms of physiology or behavior based on stress, incorporating components such as stress proxies into predictive models may, for example, increase accuracy for how a transgenic salmon will respond to environmental conditions and thus improve our understanding of the risks transgenic salmon might pose to wild populations. Interestingly, DNA methylation appeared to be independently regulated between promoter and gene regions in the liver and did not seem to play a direct role in the gene transcription responses observed in this study. While not directly influencing gene expression differences seen in the liver, this does not preclude DNA methylation as a common mechanism for gene regulation. It does suggest, however, that other gene regulatory systems, potentially originating in other organs, are responsible for gene expression differences observed in the liver. The present study has shown the remarkable responses in gene transcription that arise from environmental, genetic, and GxE influences, and has revealed underlying complex genomic mechanisms affecting plasticity and GxE interactions influencing both morphological and physiological phenotypes.

R.D. designed the experiment. J.L.L. and K.C. analyzed the data and wrote the paper with L.B., M.C., E.R., and R.D. M.C. and R.D. designed and conducted the sampling. M.C. conducted the qPCR analyses. All co-authors contributed substantially to revisions.

Funding

The Canadian Regulatory System for Biotechnology provided support for this project to R.H.D and K.A.C. Support for J.L.L. was partially provided by EPIC4 (Enhanced Production in Coho: Culture, Community, Catch) funded through Genome Canada, Genome

British Columbia, and Genome Québec with contributions from Fisheries and Oceans Canada.

Conflicts of interest: None declared.

Acknowledgments

The authors thank C. Babin, C. Hernandez, J. Létourneau and A. Perrault-Payette for conducting the bisulfite DNA libraries preparation. They thank B. Watson and H. Seshadri for sampling assistance and the fish culturists at West Vancouver Laboratory for their fish husbandry assistance. They thank the McGill University and Genome Québec Innovation Centre for their services in preparing the RNA-seq and sequencing libraries and performing the sequencing. They thank C. Rougeux, M. Laporte, Q. Rougemont and J. S. Leong for valuable discussion and comments.

Literature cited

- Alzaid A, Kim J-H, Devlin RH, Martin SAM, Macqueen DJ. 2018. Growth hormone transgenesis in coho salmon disrupts muscle immune function impacting cross-talk with growth systems. *J Exp Biol.* 221:jeb173146.
- Andrali SS, Sampley ML, Vanderford NL, Ozcan S. 2008. Glucose regulation of insulin gene expression in pancreatic beta-cells. *Biochem J.* 415:1–10.
- Baerwald MR, Meek MH, Stephens MR, Nagarajan RP, Goodbla AM, et al. 2016. Migration-related phenotypic divergence is associated with epigenetic modifications in rainbow trout. *Mol Ecol.* 25: 1785–1800.
- Björnsson BT, Johannsson V, Benedet S, Einarsdottir IE, Hildahl J, et al. 2002. Growth hormone endocrinology of salmonids: regulatory mechanisms and mode of action. *Fish Physiol Biochem.* 27: 227–242.
- Cavalli G, Heard E. 2019. Advances in epigenetics link genetics to the environment and disease. *Nature.* 571:489–499.
- Charlesworth D, Barton NH, Charlesworth B. 2017. The sources of adaptive variation. *Proc. Biol Sci.* 284:20162864.
- Cunningham CB, Ji L, McKinney EC, Benowitz KM, Schmitz RJ, et al. 2019. Changes of gene expression but not cytosine methylation are associated with male parental care reflecting behavioural state, social context and individual flexibility. *J Exp Biol.* 222: jeb188649.
- Dehkhoda F, Lee CMM, Medina J, Brooks AJ. 2018. The Growth Hormone Receptor: Mechanism of Receptor Activation, Cell Signaling, and Physiological Aspects. *Front Endocrinol.* 9:35.
- Devlin RH, Sakhrani D, Tymchuk WE, Rise ML, Goh B. 2009. Domestication and growth hormone transgenesis cause similar changes in gene expression in coho salmon (*Oncorhynchus kisutch*). *Proc Natl Acad Sci USA.* 106:3047–3052.
- Devlin RH, Sundström LF, Leggatt RA. 2015. Assessing ecological and evolutionary consequences of growth-accelerated genetically engineered fishes. *BioScience.* 65:685–700.
- Devlin RH, Yesaki TY, Biagi CA, Donaldson EM, Swanson P, et al. 1994. Extraordinary salmon growth. *Nature.* 371:209–210.
- Greenberg MVC, Bourc'his D. 2019. The diverse roles of DNA methylation in mammalian development and disease. *Nat Rev Mol Cell Biol.* 20:590–607.
- Gutierrez L, Bussell JD, Păcurar DI, Schwambach J, Păcurar M, et al. 2009. Phenotypic plasticity of adventitious rooting in Arabidopsis is controlled by complex regulation of auxin response factor transcripts and MicroRNA abundance. *Plant Cell.* 21:3119–3132.
- Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, et al. 2008. Persistent epigenetic differences associated with prenatal exposure to famine in humans. *Proc Natl Acad Sci USA.* 105: 17046–17049.
- Herman JJ, Sultan SE. 2016. DNA methylation mediates genetic variation for adaptive transgenerational plasticity. *Proc R Soc B.* 283: 20160988.
- Herrera CM, Pozo MI, Bazaga P. 2012. Jack of all nectars, master of most: DNA methylation and the epigenetic basis of niche width in a flower-living yeast. *Mol Ecol.* 21:2602–2616.
- Higgs DA, Sutton JN, Kim H, Oakes JD, Smith J, et al. 2009. Influence of dietary concentrations of protein, lipid and carbohydrate on growth, protein and energy utilization, body composition, and plasma titres of growth hormone and insulin-like growth factor-1 in non-transgenic and growth hormone transgenic coho salmon, *Oncorhynchus kisutch* (Walbaum). *Aquaculture.* 286: 127–137.
- Holm J, Møller Chr D. 1984. Growth and prey selection by Atlantic salmon yearlings reared on live freshwater zooplankton. *Aquaculture.* 43:401–412.
- Ignatz EH, Braden LM, Benfey TJ, Caballero-Solares A, Hori TS, et al. 2020. Impact of rearing temperature on the innate antiviral immune response of growth hormone transgenic female triploid Atlantic salmon (*Salmo salar*). *Fish Shellfish Immunol.* 97: 656–668.
- Kim J-H, Macqueen DJ, Winton JR, Hansen JD, Park H, et al. 2019. Effect of growth rate on transcriptomic responses to immune stimulation in wild-type, domesticated, and GH-transgenic coho salmon. *BMC Genomics.* 20:1024.
- Kronholm I, Johannesson H, Ketola T. 2016. Epigenetic control of phenotypic plasticity in the filamentous fungus *Neurospora crassa*. *G3 Genes Genomes Genet.* 6:4009–4022.
- Kuo T, McQueen A, Chen T-C, Wang J-C. 2015. Regulation of glucose homeostasis by glucocorticoids. *Adv Exp Med Biol.* 872:99–126.
- Laland K, Uller T, Feldman M, Sterelny K, Müller GB, et al. 2014. Does evolutionary theory need a rethink? *Nat. News.* 514:161–164.
- Lempiäinen H, Shore D. 2009. Growth control and ribosome biogenesis. *Curr Opin Cell Biol.* 21:855–863.
- Levis NA, Pfennig DW. 2016. Evaluating “Plasticity-First” evolution in nature: key criteria and empirical approaches. *Trends Ecol E.* 31: 563–574.
- Löhmus M, Sundström LF, Björklund M, Devlin RH. 2010. Genotype-temperature interaction in the regulation of development, growth, and morphometrics in wild-type, and growth-hormone transgenic coho salmon. *PLoS One.* 5:e9980.
- Luyer JL, Laporte M, Beacham TD, Kaukinen KH, Withler RE, et al. 2017. Parallel epigenetic modifications induced by hatchery rearing in a Pacific salmon. *Proc Natl Acad Sci USA.* 114:12964–12969.
- Martin EM, Fry RC. 2018. Environmental influences on the epigenome: exposure-associated DNA methylation in human populations. *Annu Rev Public Health.* 39:309–333.
- Mulder HA. 2016. Genomic selection improves response to selection in resilience by exploiting genotype by environment interactions. *Front Genet.* 7:178.
- Natri HM, Bobowik KS, Kusuma P, Darusallam CC, Jacobs GS, et al. 2020. Genome-wide DNA methylation and gene expression patterns reflect genetic ancestry and environmental differences across the Indonesian archipelago. *PLoS Genet.* 16:e1008749.
- Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, et al. 2010. Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.* 15:684–692.

- Oostra V, Saastamoinen M, Zwaan BJ, Wheat CW. 2018. Strong phenotypic plasticity limits potential for evolutionary responses to climate change. *Nat. Commun.* 9:1–11.
- Pacis A, Mailhot-Léonard F, Tailleux L, Randolph HE, Yotova V, et al. 2019. Gene activation precedes DNA demethylation in response to infection in human dendritic cells. *Proc Natl Acad Sci USA.* 116:6938–6943.
- Panserat S, Kamalam BS, Fournier J, Plagnes-Juan E, Woodward K, et al. 2014. Glucose metabolic gene expression in growth hormone transgenic coho salmon. *Comp Biochem Physiol A Mol Integr Physiol.* 170:38–45.
- Phillips RB, Devlin RH. 2010. Integration of growth hormone gene constructs in transgenic strains of coho salmon (*Oncorhynchus kisutch*) at centromeric or telomeric sites. *Genome.* 53:79–82.
- Raffaello A, Milan G, Masiero E, Carnio S, Lee D, et al. 2010. JunB transcription factor maintains skeletal muscle mass and promotes hypertrophy. *J Cell Biol.* 191:101–113.
- Rise ML, Douglas SE, Sakhrani D, Williams J, Ewart KV, et al. 2006. Multiple microarray platforms utilized for hepatic gene expression profiling of GH transgenic coho salmon with and without ration restriction. *J Mol Endocrinol.* 37:259–282.
- Schlichting CD, Wund MA. 2014. Phenotypic plasticity and epigenetic marking: an assessment of evidence for genetic accommodation. *Evolution.* 68:656–672.
- Smith EN, Kruglyak L. 2008. Gene–environment interaction in yeast gene expression. *PLoS Biol.* 6:e83.
- Sundström LF, Löhmus M, Devlin RH. 2016. Gene–environment interactions influence feeding and anti-predator behavior in wild and transgenic coho salmon. *Ecol Appl.* 26:67–76.
- Sundström LF, Löhmus M, Tymchuk WE, Devlin RH. 2007. Gene–environment interactions influence ecological consequences of transgenic animals. *Proc Natl Acad Sci USA.* 104:3889–3894.
- Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One.* 6:e21800.
- Thorgersen EB, Barratt-Due A, Haugaa H, Harboe M, Pischke SE, et al. 2019. The role of complement in liver injury, regeneration, and transplantation. *Hepatology.* 70:725–736.
- Van Weerd JH, Komen J. 1998. The effects of chronic stress on growth in fish: a critical appraisal. *Comp Biochem Physiol A Mol Integr Physiol.* 120:107–112.
- Vandersteen WE, Leggett R, Sundström LF, Devlin RH. 2019. Importance of experimental environmental conditions in estimating risks and associated uncertainty of transgenic fish prior to entry into. *Nat Sci Rep.* 9:1–11.
- Waltz E. 2017. First genetically engineered salmon sold in Canada. *Nat News.* 548:148–148.
- Wund MA, Baker JA, Clancy B, Golub JL, Foster SA. 2008. A test of the “flexible stem” model of evolution: ancestral plasticity, genetic accommodation, and morphological divergence in the threespine stickleback radiation. *Am Nat.* 172:449–462.
- Yeh PJ, Price TD. 2004. Adaptive phenotypic plasticity and the successful colonization of a novel environment. *Am Nat.* 164:531–542.
- Zhang X, Yang S, Chen J, Su Z. 2019. Unraveling the regulation of hepatic gluconeogenesis. *Front Endocrinol.* 9.
- Ziller MJ, Hansen KD, Meissner A, Aryee MJ. 2015. Coverage recommendations for methylation analysis by whole genome bisulfite sequencing. *Nat Methods.* 12:230–232.

Communicating editor: B. J. Andrews